

Studies on Metalloporphyrin Isomers: Synthesis, Structures, and Properties of Metal Complexes of N-Confused/N-Linked Porphyrins

マウリヤ ヨゲシュ クマール

<https://hdl.handle.net/2324/1959110>

出版情報 : Kyushu University, 2018, 博士 (工学) , 課程博士

バージョン :

権利関係 : Public access to the fulltext file is restricted for unavoidable reason (3)



氏 名 : マウリヤ ヨゲシュ クマール

論 文 名 : Studies on Metalloporphyrin Isomers: Synthesis, Structures, and Properties of Metal Complexes of N-Confused/N-Linked Porphyrins (金属コロール異性体に関する研究 : N-混乱/N-連結コロール金属錯体の合成、構造および物性)

区 分 : 甲

論 文 内 容 の 要 旨

Corroles, one of the important family of porphyrinoids, are contracted tetrapyrrolic macrocycles containing a direct pyrrole-pyrrole linkage that resemble the skeletal structure of corrin. The characteristic feature of corroles is an ability to coordinate metal ions with unusual higher oxidation states, which allowed us to apply to catalysts. To date, researchers have made tremendous efforts to elucidate the structure-property relationships of the metallo-corroles. However, the studies of bona fide metalloporphyrin isomers are still in its infancy in analogy with the porphyrin derivatives.

In this thesis, the author described the wide scope of metalloporphyrin analog chemistry, that is N-confused/N-linked corroles and their metal complexes. The intriguing photophysical properties and redox reactivities of the metal complexes of these corrole mutants have been investigated, which arose from the special coordination environment with an inner carbon donor in the confused pyrrolic ring of the respective macrocycles. The findings presented in this thesis would provide an important basis for the future applications, such as sensors, emitters, and catalysts.

Chapter 1 outlines a brief overview of the chemistry of porphyrinoids, mainly focusing on the fundamental structural properties, reactivities, and the potential application of corroles and metalloporphyrins. The important question why the high-valent metal corrole complexes can be a competent example for a wide variety of applications has been discussed, while a combination of some specific metal ions with redox-active corrole ligands gave rise to a varied degree of ligand noninnocence character. In this aspect, core modifications of the corrole skeleton with “N-confusion” approach can provide a new redox chemistry for carbacorroles and their metal complexes toward unique catalytic activities. This chapter also shed light

on the class of covalently linked corrole dimers which can be considered as an example for biomimetic models and can be promising candidates for optoelectronic devices using the distinct near-infrared phosphorescent properties.

In Chapter 2, a concept of symmetry modulation of π -HOMO (b1) orbital of the corroles by N-confusion approach was implemented to stabilize the distinct high valent copper species using the aforementioned NNNC coordination environment with a strong σ donating ability of the inner carbon atom. This peculiar core-modified corrole framework indeed enabled to stabilize the high valent organocopper(III) species predominantly as inferred from the complementary spectroscopic means; X-ray crystallography, variable temperature NMR, XPS, XANES, and UV-vis absorption. The meticulous tuning of the orbital energies of Cu $d_{x^2-y^2}$ and the π -carbacorrole (b1) debilitates its interaction in the electronic structures, and the ground singlet Cu(III) structures were thus verified by DFT calculations. Furthermore, these organocopper corroles demonstrated unique redox reactivity to give a new π -laminated dimeric architecture via an oxidative inner carbon-carbon bond formation adjacent Cu(III) center.

In Chapter 3, utilizing a facile synthetic strategy for covalently linked corrole dimers that the author established, a series of conformationally stable chiral chromophores have been developed based on the novel corrole isomers. These face-to-face dimeric corroles possessing different ground electronic structures display unique chirality structures (e.g., stereogenic centers and axial chirality) with extraordinary high geometric stabilities, which can be regarded as mimetic model platforms for photosynthetic reaction centers, dimeric chlorophylls.

In Chapter 4, the author describes the synthesis of a series of iridium(III) complexes of the corrole analogs possessing various *para*-substituted pyridine axial ligands as novel near-infrared emitting materials at wavelength beyond 900 nm. The comparative studies have been made with the iridium corrole congeners to understand how the N-linked mutation affects the photophysical properties. Moreover, various axial pyridine ligands on the complexes were utilized to control the inherent stability and solubility in organic solvents and subsequently can tune their photophysical properties.

In Chapter 5, the author summarized the findings and perspectives of the future works. These metal complexes of carbacorrole analogs obtained by the N-/Neo-confusion modification can be utilized for various applications in catalysts, chiral recognition sensors and NIR-responsive devices.

〔作成要領〕

1. 用紙はA4判上質紙を使用すること。
2. 原則として、文字サイズ10.5ポイントとする。
3. 左右2センチ，上下2.5センチ程度をあげ，ページ数は記入しないこと。
4. 要旨は2,000字程度にまとめること。
(英文の場合は，2ページ以内にまとめること。)

5. 図表・図式等は随意に使用のこと。
6. ワードプロセッサで書すること（手書きする場合は楷書体）。

この様式で提出された書類は，「九州大学博士学位論文内容の要旨及び審査結果の要旨」の原稿として写真印刷するので，鮮明な原稿をクリップ止めで提出すること。