

Effect of brisk walking on the body weight, body fat, waist circumference and blood lipids in adults: A systematic review

ZHANG, Tengfei

Graduate School of Human-Environment Studies, Kyushu University

CHEN, Si

Graduate School of Human-Environment Studies, Kyushu University

SAITO, Atsushi

Faculty of Human-Environment Studies, Kyushu University

<https://doi.org/10.15017/2230692>

出版情報：健康科学. 41, pp.23-39, 2019-03-20. 九州大学健康科学編集委員会
バージョン：
権利関係：



—Review—

Effect of brisk walking on the body weight, body fat, waist
circumference and blood lipids in adults:
A systematic review

Tengfei ZHANG¹⁾, Si CHEN¹⁾, and Atsushi SAITO²⁾*

Abstract

Purpose: The purpose of this review was to assess the effect of brisk walking on the body weight, body fat, waist circumference and blood lipid values in adults.

Methods: In August 2018, we conducted electronic searches of PubMed and Web of Science to identify pertinent studies; those with an experiment period exceeding two weeks, human randomized controlled trials (RCTs), and those that only assessed brisk walking was included.

Results: A total of 523 citations were identified from our search of the electronic literature, and 29 RCTs that included 1697 individuals were identified. Overall, 15 studies found a significant change in the decrease in body weight, body fat, and waist circumference. In addition, eight of these studies demonstrated statistically significant improvements in the low density lipoprotein cholesterol (LDL), high-density lipoprotein cholesterol (HDL), triglyceride (TG) and total cholesterol (TC) values. **Conclusion:** Brisk walking was able to reduce the body weight, body fat and waist circumference, and improved the levels of blood lipids. However, these effects are associated with the amount and intensity of exercise and the state of health of the participants.

Key words: brisk walking, adults, blood lipids, body weight, body fat

(Journal of Health Science, Kyushu University, 41: 23-39, 2019)

1 Graduate School of Human-Environment Studies, Kyushu University, Fukuoka, Japan.

2 Faculty of Human-Environment Studies, Kyushu University, Fukuoka, Japan.

*Correspondence to: Faculty of Human-Environment Studies, Kyushu University, 744 Motoooka, Nishi-ku, Fukuoka City 819-0395, Japan.

Tel: +81-92-802-5159 E-mail: saito-a@ihs.kyushu-u.ac.jp

Introduction

It is widely assumed that regular physical activity can help prevent primary and secondary chronic diseases¹⁾. Most countries' government and health organization recommend that subjects engage in regular moderate or vigorous intensity physical activity. However, modern society has changed our approach to work, and the modern lifestyle has also changed our behaviors. Whether working or enjoying leisure time, we spend a great proportion of our life engaged in sedentary behavior. Sedentary behavior is defined as any waking behavior characterized by an energy expenditure (as well as postural 「sitting or reclining」 and contextual 「waking」) ≤ 1.5 metabolic equivalents (METs)²⁾. According to this definition, a person may be described as sedentary if they engage in a large amount of sedentary behavior. However, there was different energy expenditure in sedentary. For example, sitting in a classroom is assigned 1.8METs, while driving a car is assigned 2.5METs. There is no consistent definition of sedentary behavior; nevertheless, a common phenomenon exist namely a lack of ambulatory movement in any posture. This is an important characteristic that distinguishes sedentary behavior from physical activity. Physical activity is defined as any body movement generated by the contraction of skeletal muscles, such activity should increase energy expenditure above the resting metabolic rate. Physical activity is characterized by its modality, frequency, intensity, duration, and context of practice³⁾. A number of studies have found moderate-to-strong evidence that diabetes and mortality are associated with sedentary behavior^{4,5)}. Warren et al.⁶⁾ found that sedentary behavior and riding in a car were significant predictors of mortality due to cardiovascular disease (CVD). A review study showed that

increased physical activity helps prevent chronic disease and promote public health⁷⁾.

Sedentary behavior is a major factor that increases the risk of chronic diseases, and it also significantly affects on one's body weight. Over the last three to four decades, the prevalence of overweight and obesity has increased dramatically. For instance, in 2015, about 107.7 million children and 603.7 million adults were obese around the world. Obesity is a major risk factor for CVD, type 2 diabetes mellitus (T2DM), hypertension, coronary artery diseases, metabolic syndrome, and some cancers⁸⁾. It is also an important risk factor for death due to non-communicable disease.

Over nutrition is an obvious candidate contributor to the rise in obesity, but increased rates of physical inactivity and sedentary behavior are also implicated. Several studies have found that regular moderate physical activity was able to improve blood lipids, reduce the risk of CVD diseases, and prevent and control T2DM. It is also a good method of managing obesity. The World Health Organization⁹⁾ recommends that adults spend at least 150 minutes per week engaged in aerobic physical activity (moderate intensity) per week, or at least 75 minutes of high-intensity physical activity. The American College of Sports Medicine has similar recommendations in its physical activity guidelines.

Walking has been described as a near perfect "exercise" and has been recommended by health organizations worldwide¹⁰⁾. Walking programs have a higher level of adherence than other forms of exercise¹¹⁾. Brisk walking as a simple, economic and safe form of exercise is the most popular moderate-intensity aerobic physical activity for both men and women, especially for most middle-aged and/or overweight and obesity individuals. It is the most likely exercise or

physical activity to be chosen to improve chronic diseases, and T2DM and to manage body weight¹²). According to the definition of moderate-intensity activity, walking is “brisk” at a speed of 2.5 to 4.2 mph on firm ground. Studies have shown that 30 minutes of moderate-to-vigorous walking corresponds to a total of 3100-4000 steps¹³). Previous studies have demonstrated that brisk walking can result in improved low density lipoprotein cholesterol (LDL-C), total cholesterol (TC)¹⁴, and high-density lipoprotein cholesterol (HDL-C)¹⁵). However, conflicting results have also been reported, with some studies showing that brisk walking has no substantial effect on the blood lipid values^{16,17}), although a systematic review and meta-analysis showed that brisk walking was able to significantly reduce the body weight, body mass index (BMI), waist circumference, and fat mass in obese subjects¹⁸). Brisk walking proved to be an economic and feasible form of exercise that can be performed by nearly all individuals (especially for those with overweight and obesity), in all kinds of places and at almost any time.

The aim of this system review was to systematically summarize, analyze, and interpret the health benefits of brisk walking in human with regard to its effects on body weight, the BMI, waist circumference, HDL, LDL, TC, triglyceride (TG), and other health-related measures.

Method

Literature search strategy

This systematic review followed the Cochrane systematic review guidelines. The online database Web of Science and PubMed were evaluated through August 3, 2018, for relevant periodicals. The PICO patient, intervention, comparison, and outcome (PICO) approach was used in this review search strategy. Our search eliminated two sets of

keywords using the Boolean operator “NOT”, and combined two sets of keywords using “AND”, with “OR” used to combine the keywords. Our keywords search strategy included both full and abbreviated terms as follows: brisk walking, adults, body weight, waist circumference, BMI, LDL-C, TC, HDL-C, fat ratio, and TG. The search initially yielded 523 potentially relevant articles. The duplicate articles were removed based on titles and abstracts. Due to the number of results, a second researcher independently screened a random sample of 20% of the retrieved record, and completed the inclusion of articles (Figure 1).

Inclusion and exclusion criteria

A study was included if the following requirements were met: (a) subjects ≥ 18 years old; (b) randomized trial; and (c) brisk walking or moderate exercise evaluated. The exclusion criteria were as follows: (a) study period of less than 2 weeks; (b) subjects ≤ 18 years; (c) no relevant results; (d) a Nordic walking study or walking program study (not brisk walking).

Assessing the quality

The quality of the studies was assessed using the Cochrane Risk of Bias Tool. The criteria included random allocation, blinding of supplement, outcome assessors and individuals, allocation concealment, use of an intention-to-treat analysis, and other biases. Two researchers assessed the studies, separately. If any differences in their findings were noted, the conflict was resolved by a discussion.

Data extraction

Two researchers extracted all relevant data independently. One author extracted the following data from the eligible articles: the first author and publication year, study design, intervention duration, participants, intervention characteristics, control group characteristics, exercise (volume and intensity),




Figure 1 Flow diagram showing the literature search and selection process

and outcomes of body weight, waist circumference, BMI, LDL-C, TC, HDL-C, TG.

Results

A total of 523 RCTs studies were identified through initial searches. Following the removal of duplicates, titles, and abstracts were screened, resulting in 72 articles undergoing full-text screening for inclusion. Overall, 29 RCT articles were identified (Fig. 1). We excluded 34 articles for the following reasons: 6 were testing studies; 17 were combined with other exercises; 8 did not include indices, and 3 had no sedentary control group. This review identified a total of 1697 individuals (brisk walking group, $n=931$; control group, $n=766$), investigating the long-term health effects of brisk walking.

Mean research duration was 3-48 weeks. The health of participants varied among studies, including overweight and/or obese, T2DM, nonalcoholic fatty liver, physically deconditioned, hypercholesterolemic, and eumenorrheic patients. Two studies did not report the values of body weight, body fat or waist circumference. And nine did not report the values of LDL, HDL, TG or TC (Table 1).

Brisk walking intensity

The definition of brisk walking differed among the studies included in this review. Regarding the definition of brisk walking, the maximum heart rate ranged from 40% to 80%. Several studies were based on breathing difficulty, faster than normal, $62 \pm 2\%$ VO_2max , 3~5 km/h. In addition, one study did not clearly define the meaning of brisk walking used in their trial. Eight studies reported that

the definition of brisk walking was based on the participants' self reported description.

Body weight and waist circumference

Four studies did not report the body weight. And 13 did not report waist circumference. Twelve studies included only women participants, 3 included only men, and 14 included both men and women participants. Thirteen studies^{26,27,28,30,33,38,39,40,42,46,47,48,50} found a significant decrease in the body weight at the end of the study period. Twelve studies failed to show a significant decrease in the body weight, but seven of these trails included both men and women as participants; the physiology differs between sexes, which may have been a major confounding factor in our review. John's study, the first evaluated in this review, found that brisk walking significantly reduced the body weight in the intervention group compared with the control group. This was a long-term study, and the participants were sedentary obese women, however, the participants' BMI was not reported. Another study¹⁹) showed that sedentary obesity and a long intervention period were important factors influencing the reduction in body weight.

Body fat

Seventeen trials measured the effect of brisk walking on body fat. Eight of these studies^{26,27,38,39,40,46,50,52}) showed statistically significant decreases in body fat after intervention.

LDL and TC

Seventeen trials measured the effect of brisk walking on the LDL and TC levels. Three of these trials^{28,32,42}) found statistically significant reductions in the LDL levels; however, one of these three studies included hypertensive participants, another had only a three-week intervention period and the third did not describe the BMI of the participants. Two^{40,42}) of seventeen studies shown a significant reduction in the TC levels;

however, one of these two studies included hypercholesterolemic patients, and the other only had a three-week intervention period.

HDL and TG

Nineteen trials measured the effects of brisk walking on the HDL-C levels. Four^{24,32,43,45}) of these studies found statistically significant reductions in the HDL-C levels. However, the BMI of the participants was not mentioned in two of these studies, another included hypertensive participants, and the remaining one included type 2 diabetes patients. Seventeen trials measured the effect of brisk walking on the TG levels. Only two^{37,40}) of these studies showed statistically significant reductions in TG. However, only two of these studies showed statistically significant reductions in the TG levels, with one including hypertensive subjects and the other one including high-triglyceride value subjects.

Discussion

The aim of this review was to assess the effect of brisk walking on the body weight, body fat, waist circumference, and blood lipid values in adults. A total of 29 intervention studies met the inclusion criteria and were included in this review. While all of the included trials assessed brisk walking as the modality, the intensity of the exercise varied widely. Brisk walking has been confirmed as a suitable activity for maintaining a healthy lifestyle by the world health organization, public health agencies, and health research institutions around the globe. However, the precise definition of brisk walking remains unclear. Porcari et al.²⁰) showed that brisk walking is defined as a maximal heart rate of $\geq 70\%$. Spelman et al.²¹) similarly suggested that the maximal heart rate should be above 70%. Previous studies^{22,23}) have shown that brisk walking is defined as a speed of walking ranging from three to four miles per hour.

Table 1 Basic characteristic of included 29 studies

Study	State of health	Participants	N	Age(years)	BMI	Intervention	Weeks	Intensity	physical fitness	Results blood lipids
Hardman A et al. (1989) ²⁴⁾	sedentary	women	44	44.9 ± 7.9; 44.4 ± 9.2	unknown	IG=28, 155±48min/week, BW, 16.1-17.4km/week; CG=16, habitual lifestyle	48	1.72±0.26m/s~ 1.87±0.37m/s	unknown	HDL ↑ TC: none
Sylvia RC et al. (1991) ²⁵⁾	sedentary overweight and obesity	women	35	36.0 ± 1.6; 32.4 ± 1.5	28.3 ± 0.7 27.8 ± 0.9	IG=18, 45min/week, 5 time/week, 62±2% VO _{2max} ; CG=17, habitual lifestyle	15	62±2% VO _{2max}	BW none BF: none	unknown
John PF et al. (1993) ²⁶⁾	obesity	men and women	165	25~45	unknown	IG=43, 45min/time, 3~5 sessions/week, CG=38, habitual lifestyle	48	self-monitoring; breathing difficulty.	BW ↓ BF ↓ WC ↓	unknown
Stensel DJ et al. (1993) ²⁷⁾	sedentary healthy	men	72	42~59	25.3 ± 0.4 24.9 ± 0.6	IG=48, 20~25min to 40~45min/day, 2~3 sessions/week, CG=24, habitual lifestyle	48	self-monitoring	BW ↓ BF ↓ WC :none	HDL: none TC: none LDL: none TG: none
Aldred HE et al. (1995) ²⁸⁾	sedentary (overweight/ obesity)	women	26	41~55	unknown	IG=13, 60~180min/week, 3time/week, 20~50min/session, CG=13, habitual lifestyle	12	self-monitoring	BW ↓ WC :none	HDL: none TC: none LDL: ↓ TG: none
Santiago MC et al. (1995) ²⁹⁾	sedentary eumenorrheic women	women	32	22~40	24.6 ± 3.8 23.6 ± 3.5	IG=21, 4.8km/week, 4time/week, mean intensity 72% maximal heart rate ; CG=11, habitual lifestyle	40	72% maximal heart rate	BW none BF none	HDL: none

Brooke-Wavell K et al. (1997) ³⁰⁾	sedentary post-menopausal women (health and overweight)	84	60~70	25.8± 3.8 25.6± 3.5	IG=43, 20~50min/day, 7time/week ; CG=41, habitual lifestyle	48	self-monitoring	BW ↓	unknown
Coleman KJ et al. (1999) ³¹⁾	sedentary men (health and overweight)	36	18~45	25.0± 3.4 26.0± 4.5 26.4 ±4.1	IG1=12, 30min/day, 6time/week; IG2=12, 3×10min/day, 6 time/week; CG=12, self walk;	16	accelerometer	BW none	unknown
Higashi Y et al. (1999) ³²⁾	Hypertensive Subjects	27	53±10; 51±8	24.0± 1.7 24.2± 1.8	IG1=20, 30min/day, 5~7 time/week; CG=7, habitual lifestyle;	12	self-monitoring; 52 ± 6%	BW none	HDL: ↑ TC: none LDL: ↓ TG: none
Brooke-Wavell K et al. (2001) ³³⁾	Postmenopausal Women (overweight/ obesity)	68	60~70	unknown	IG=16, 120~280min/ two weeks, 3time/week, 20~45min/session ; CG=20, habitual lifestyle	48		BW ↓	unknown
Murtagh EM et al. (2005) ³⁴⁾	health	30	45.7± 9.4	unknown	IG=8, 20min/day, 3 time/week; CG=7, habitual lifestyle;	12	heart rate; treadmill	BW : none BF :none WC : none	HDL: none TC: none LDL: none TG: none

Tully MA et al. (2005) ³⁵⁾	sedentary, overweight and obesity	men and women	31	50~65	28.09± 3.98 27.29 ±3.25	IG=17, 30min/day, 3 time/week; CG=9, habitual lifestyle;	12	self-monitoring	BW : none BF : none WC : none	HDL: none TC: none LDL: none TG: none
Audette JF et al. (2006) ³⁶⁾	sedentary; (health and overweight)	elderly women	27	71.4 ± 4.5	unknown	IG=10, 60min/day, 3 time/week; CG=8, habitual lifestyle;	12	maximal heart rate: 50%~70%	unknown	unknown
Lee MR et al. (2006) ³⁷⁾	overweight and obesity; high triglyceride	women	19	35 ~ 50	26.25± 1.83 26.53 ±1.38	IG=9, 20~50min/day, 3 ~6 time/week; CG=10, BW plus diet;	12	maximal heart rate: 40-50% ~ 60-70%	unknown	TG ↓
Kirkwood L et al. (2007) ³⁸⁾	overweight and obesity	women	69	30 ~ 50	31.6 ±3.8 31.6± 4.2	IG=19, 60min/day, 6 time/week; CG=18, habitual lifestyle;	12	accelerator monitors	BW ↓ BF ↓ WC ↓	HDL: none TC: none LDL: none TG: none
Tully MA et al. (2007) ³⁹⁾	sedentary (overweight and obesity)	men and women	106	40 ~ 61	27.79± 4.85 25.77± 4.09 26.32 ±5.94	IG1 =44, 30min/day, 3 time/week; IG2 =42, 30min/day, 5 time/week; CG=20, habitual lifestyle;	12	faster than normal, pedometer	BW ↓ BF ↓ WC ↓	HDL: none TC: none LDL: none TG: none

Coghill N et al. (2008) ⁽⁴⁰⁾	hypercholesterolaemic (overweight and obesity)	men	67	55.1 ± 4.9	28.13 ± 3.86 27.27 ± 2.98	IG = 38, 30min/day, 5 - 7 time/week; CG=29, habitual lifestyle;	12 accelerator monitors; RPE: 12-14	BW ↓ BF ↓ WC ↓	HDL: none TC ↓ LDL: none TG ↓
Pierce GL et al. (2011) ⁽⁴¹⁾	sedentary	men and women	22	55 - 79	25.3 ± 0.7 25.1 ± 0.8	IG = 26; 40 - 50 min/day, 6 - 7 time/week; CG=10, habitual lifestyle;	12 maximal heart rate: 70%~75%	BW : none BF : none WC : none	HDL: none TC: none LDL: none TG: none
Pagels P et al. (2012) ⁽⁴²⁾	health	men and women	33	25 - 45	26.2 ± 4.4 23.7 ± 2.3	IG = 16; 30 min/day, 7 time/week; CG=17, habitual lifestyle;	3 pedometer and accelerometer	BW ↓	HDL: none TC ↓ LDL ↓ TG: none
Foulds,HJ et al. (2014) ⁽⁴³⁾	health and overweight	men and women	72	44 ± 13.0	unknown	IG 1 = 10; 10min/week; IG 2 = 10; 10min/day, 3 time/week; IG 3 = 18; 30min/day, 3 time/week; IG 4 = 10; 60min/day, 3 time/week; CG=10, habitual lifestyle;	13 self-report	unknown	HDL ↑ IG2; IG4 TC: none LDL: none TG: none
Kearney TM et al. (2014) ⁽⁴⁴⁾	sedentary; overweight and obesity	men and women	77	30 - 55	29.3 ± 4.3 29.2 ± 4.3	IG = 16; 3 × 10 min/day, 5 time/week; CG=17, habitual lifestyle;	24 self-report	BW : none BF : none WC : none	HDL: none TC: none LDL: none TG: none

Johnson ST et al. (2015) ⁽⁴⁵⁾	Type 2 Diabetes	postmeno pausal women	198	59.5 ± 8.3	34.6 ± 6.4 32.5 ± 6.4	IG = 102; 30 min/day, 3 time/week; CG=96, habitual lifestyle;	24	self-monitoring; pedometer	BW : none WC : none	HDL ↑ LDL: none
Hui SS et al. (2016) ⁽⁴⁶⁾	no exercise habits	men and women	374	36 -60	23.2± 3.0 22.7 ± 3.3	IG = 121; 45 min/day, 5 time/week; CG=124, habitual lifestyle;	12	self-monitoring	BW : ↓ BF ↓	unknown
Zhang HJ et al. (2016) ⁽⁴⁷⁾	Nonalcoholic fatty liver	men and women	220	53.9 ± 7.1	28.1 ± 3.3 28.0± 2.7	IG=73, 150min/week; CG=74, habitual lifestyle;	48	maximal heart rate: 45%~55%; pedometer	BW ↓ BF : none WC ↓	HDL: none TC: none LDL: none TG: none
Blain H et al. (2017) ⁽⁴⁸⁾	sedentary and physically deconditioned	women	121	65.7 ± 4.3	24.25± 3.46 26.64 ± 4.22	IG=51, 150min/week, 3 time/week; CG=47, habitual lifestyle;	24	maximal heart rate: 40%~80%; HR monitor	BW ↓ BF : none	unknown
Buyukyazi G et al. (2017) ⁽⁴⁹⁾	health	premenop ausal women	30	33 - 50	23.6 ~ 29.6 20.9 ~ 30.1	IG=11, 30 - 51 min/week, 3 time/week; CG=8, habitual lifestyle;	8	maximal heart rate: 70%~75%; HR monitor	BW : none WC : none	unknown
Herzig KH et al. (2014) ⁽⁵⁰⁾	health	men and women	78	58.8±10.4	32.6± 5.7 30.9 ± 4.8	IG=33, 60 min/week, 3 time/week; CG=35, habitual lifestyle;	12	3 ~ 5 km/h	BW ↓ BF ↓ WC ↓	HDL: none TC: none LDL: none TG: none

Ho SS et al. (2012) ⁽⁵¹⁾	overweight and obesity	men and women	81	40 - 66	32.7± 1.3 32.4 ± 1.4	IG=15, 30 min/week, 5 time/week; CG=16, habitual lifestyle;	12	maximal heart rate: 60%±10%; treadmill	BW : none BF : none WC ↓	HDL: none TC: none LDL: none TG: none
McNeilly AM et al. (2011) ⁽⁵²⁾	sedentary; obesity	men and women	24	54 ± 8	33 ± 6 32± 7	IG=12, 30 min/week, 5 time/week; CG=12, habitual lifestyle;	12	maximal heart rate: 65%; monitor	BW : none BF ↓ WC ↓	HDL: none TC: none LDL: none TG: none

N = number; BMI = body mass index; IG = intervention group; CG = control group; BW = body weight; BF = body fat; WC = waist circumference; HDL = high-density lipoprotein cholesterol; LDL = low density lipoprotein cholesterol; TC = total cholesterol; TG = triglyceride.

However, only 7 trials in our review reported that a maximal heart rate $\geq 70\%$ was achieved. Other studies used self-monitoring to assess the presence of brisk walking (accelerator, pedometer and rate of perceived exertion scale). One of these studies reported that the maximal oxygen consumption should be $62\% \pm 2\%$ for brisk walking. However, another study that assessed the speed and intensity of walkers showed that the threshold of maximal oxygen consumption was $68.6\% \pm 14.9\%$.

A systematic review and meta-analysis⁵³⁾ demonstrated that brisk walking was able to reduce the body weight, body fat and waist circumference in obese adults. Twelve studies in our review reported statistically significant changes in the body weight, and eight reported statistically significant changes in the body fat and waist circumference. The intensity of brisk walking differed markedly among our included studies. Nemoto et al.⁵⁴⁾ found that brisk walking, defined as a maximal oxygen consumption of 50%-70%, was able to reduce the body mass. Blain et al.⁴⁸⁾ also showed that brisk walking, defined as a maximal oxygen consumption of 60% - 80%, was able to reduce the body mass and body fat of sedentary women. Maintaining or increasing the amount of moderate or vigorous intensity physical activity is recommended by a number of health organizations to reduce the risk of chronic diseases in sedentary adults. As the one of the most popular physical activities, brisk walking could reduction the body mass, body fat or waist circumference. However, the recommended levels of such physical activities have generally not led to clinically significant body mass loss⁵⁵⁾. Previous studies^{56,57)} also have shown that statistically significant weight loss generally does not occur with moderate-intensity physical activity. However, it was able to change the other favorable cardiometabolic. Thus, while brisk

walking can reduce body weight, no significant reduction in body weight may occur.

Several studies have shown that physical activity has favorable effects on the blood lipid and lipoprotein profiles. A previous review⁵⁸⁾ found that exercise increased the HDL-C levels and reduced those of TG and LDL. However, other studies on the effects of aerobic exercise on the lipids and lipoproteins in adults with cardiovascular disease have yielded conflicting results, with the changes in the blood lipid values not always being statistically significant. Indeed, only six trials among the studies included in our review reported statistically significant reductions in the blood lipid and lipoprotein values, and four of these studies included chronic disease patients as participants. Another meta-analysis showed that aerobic exercise increased the HDL and decreased the TG levels in adults, especially men, with cardiovascular disease⁵⁹⁾. This finding, along with others, confirmed that physical activity could improve the blood lipid and lipoprotein levels. Physical activity can not only improve the health of chronic disease patients, but also reduce the risk of chronic disease altogether in healthy humans. A meta-analysis found that the effect of physical activity on blood lipids was associated with increased levels of such activity, and Duncan et al.⁶⁰⁾ found that a difference in the amount of-exercise appears to result in a different outcome in the blood lipid values. Another study⁶¹⁾ showed that different intensity levels also significantly affected the lipoprotein levels.

In conclusion, present review found that a brisk walking program was able to reduce the body weight, body fat, and waist circumference, as well as reduce the levels of LDL, TC and TG, and increase the levels of HDL. Furthermore, two factors were found to influence the effects of brisk walking on the

body weight, body fat, waist circumference and blood lipids: (1) the amount and intensity of the brisk walking, and (2) the state of health of the participants.

Limitation

Several limitations associated with the present review warrant mention. First, our review was confined to published research, and we simply searched PubMed and Web of Science to find appropriate articles. Second, there may have been some bias in a few of the selected studies, which may have influenced the results. Finally, the health of the subjects among the different studies may have influenced the findings, including type 2 diabetes mellitus, physically deconditioned, hypercholesterolemic, and eumenorrheic patients. Such limitations may thus have been major confounding factors in this review.

References

- 1) Warburton DE, Nicol CW, Bredin SS (2006): Health benefits of physical activity: the evidence. *CMAJ*, 174: 801–809.
- 2) Sedentary Behaviour Research Network (2012): Letter to the editor: standardized use of the terms ‘sedentary’ and ‘sedentary behaviours’. *Appl Physiol Nutr Metab*, 37: 540–542.
- 3) Thivel D, Tremblay A, Genin PM, Panahi S, Rivière D, Duclos M (2018): Physical Activity, Inactivity, and Sedentary Behaviors: Definitions and Implications in Occupational Health. *Front Public Health*, 6:1–5.
- 4) Van Uffelen JGZ, Wong J, Chau JY, Ploeg HP, Riphagen R, Gilson ND, Burton NW, Healy GN, Thorp AA, Clark BK, Gardiner PA, Dunstan DW, Bauman A, Owen N, Brown WJ (2010): Occupational sitting and health risks: a systematic review. *Am J Prev Med*, 39:379–388.
- 5) Proper KI, Singh AS, van Mechelen W, Chinapaw MJM (2011): Sedentary behaviors and health outcomes among adults: a systematic review of prospective studies. *Am J Prev Med*, 40:174–182.
- 6) Warren TY, Barry V, Hooker SP, Sui X, Church TS, Blair SN (2010): Sedentary behaviors increase risk of cardiovascular disease mortality in men. *Med Sci Sports Exerc*, 42:879–85.
- 7) Haskell WL, Blair SN, Hill JO (2009): Physical activity: health outcomes and importance for public health policy. *Prev Med*, 49:280–282.
- 8) Malik VS, Willett WC, Hu FB (2013): Global obesity: trends, risk factors and policy implications. *Nat Rev Endocrinol*, 9:13–27.
- 9) World Health Organization (2010): Global recommendations on physical activity for health. Geneva: World Health Organization.
- 10) Morris J, Hardman A (1997): Walking to health. *Sports Med*, 23: 306–332.
- 11) Siegel PZ, Brackbill RM, Heath GW (1995): The epidemiology of walking for exercise: implications for promoting activity among sedentary groups. *Am J Public Health*, 85:706–771.
- 12) Lee IM, Buchner DM (2008): The importance of walking to public health. *Med Sci Sports Exerc*, 40:S512–S518.
- 13) Pillay JD, Kolbe-Alexander TL, Proper KI, van Mechelen W, Lambert EV (2014): Steps that count: physical activity recommendations, brisk walking, and steps per minute—how do they relate? *J Phys Act Health*, 11:502–508.
- 14) Pagels P, Raustorp A, Archer T, Lidman U, Alricsson M (2012): Influence of moderate, daily physical activity on body composition and blood lipid profile in Swedish adults. *J Phys Act Health*, 9:867–874.
- 15) Murphy M, Nevill A, Neville C, Biddle S, Hardman A (2002): Accumulating brisk walking for fitness, cardiovascular risk, and

- psychological health. *Med Sci Sports Exerc*, 34:1468-1474.
- 16) Murtagh EM, Boreham CA, Nevill A, Hare LG, Murphy MH (2005): The effects of 60 minutes of brisk walking per week, accumulated in two different patterns, on cardiovascular risk. *Prev Med*, 41:92-97.
 - 17) Tully MA, Cupples ME, Chan WS, McGlade K, Young IS (2005): Brisk walking, fitness, and cardiovascular risk: a randomized controlled trial in primary care. *Prev Med*, 41:622-628.
 - 18) Mabire L, Mani R, Liu L, Mulligan H, Baxter D (2017): The influence of age, sex and body mass index on the effectiveness of brisk walking for obesity management in adults: a systematic review and meta-analysis. *J Phys Act Health*, 14:389-407.
 - 19) Ohkawara K, Tanaka S, Miyachi M, Ishikawa-Takata K, Tabata I (2007): A dose-response relation between aerobic exercise and visceral fat reduction: systematic review of clinical trials. *Int J Obes*, 31: 1786-1797.
 - 20) Porcari JP, McCarron R, Kline G, Freedson PS, Ward A, Ross JA, Rippe JM (1987): Is fast walking an adequate aerobic training stimulus for 30 to 69 year-old men and women? *Phys Sports Med*, 1: 119-129.
 - 21) Spelman CC, Pate RR, Macera CA, Ward DS (1993): Self-selected exercise intensity of habitual walkers. *Med Sci Sports Exerc*, 25:1174-1179.
 - 22) Suter E, Marti B, Gutzwiller F (1994): Jogging or walking: comparison of health effects. *Ann Epidemiol*, 4:375-381.
 - 23) Pate RR, Pratt M, Blair SN, Haskell WL, Macera CA, Bouchard C, Buchner D, Ettinger W, Heath GW, King AC, Kriska A, Leon AS, Marcus BH, Morris J, Paffenbarger Jr RS, Patrick K, Pollock ML, Rippe JM, Sallis J, Wilmore JH (1995): Physical activity and public health: a recommendation from the Centers for Disease Control and Prevention and the American College of Sports Medicine. *JAMA*, 273:402-407.
 - 24) Hardman AE, Hudson A, Jones PR, Norgan NG (1989): Brisk walking and plasma high density lipoprotein cholesterol concentration in previously sedentary women. *BMJ*, 299:1204-1205.
 - 25) Cramer SR, Nieman DC, Lee JW (1991): The effects of moderate exercise training on psychological well-being and mood state in women. *J Psychosom Res*, 35: 437-449.
 - 26) John PF, Ken GG, Rebecca SR, Raynaud AS, Darnell L, Brown AH, Gotto AM (1993): Response of free-living adults to behavioral treatment of obesity: attrition and compliance to exercise. *Behavior Therapy*, 24: 659-669.
 - 27) Stensel DJ, Hardman AE, Brooke-Wavell K, Vallance D, Jones PR, Norgan NG, Winder AF (1993): Brisk walking and serum lipoprotein variables in formerly sedentary men aged 42-59 years. *Clin Sci*, 85: 701-708.
 - 28) Aldred HE, Hardman AE, Taylor S (1995): Influence of 12 weeks of training by brisk walking on postprandial lipemia and insulinemia in sedentary middle-aged women. *Metabolism*, 44: 390-397.
 - 29) Santiago MC, Leon AS, Serfass RC (1995): Failure of 40 weeks of brisk walking to alter blood lipids in normolipemic women. *Can J Appl Physiol*, 20: 417-428.
 - 30) Brooke-Wavell K, Jones PR, Hardman AE (1997): Brisk walking reduces calcaneal bone loss in post-menopausal women. *Clin Sci*, 92: 75-780.
 - 31) Coleman KJ, Raynor HR, Mueller DM, Cerny FJ, Dorn JM, Epstein LH (1999): Providing sedentary adults with choices for meeting their walking goals. *Prev Med*, 28: 510-519.

- 32) Higashi Y, Sasaki S, Sasaki N, Nakagawa K, Ueda T, Yoshimizu A, Kurisu S, Matsuura H, Kajiyama G, Oshima T (1999): Daily aerobic exercise improves reactive hyperemia in patients with essential hypertension. *Hypertension*, 33: 591-597.
- 33) Brooke-Wavell K, Jones PR, Hardman AE, Tsuritan, Yamada Y (2001): Commencing, continuing and stopping brisk walking: effects on bone mineral density, quantitative ultrasound of bone and markers of bone metabolism in postmenopausal women. *Osteoporos Int*, 12: 581-587.
- 34) Murtagh EM, Boreham CA, Nevill A, Hare LG, Murphy MH (2005): The effects of 60 minutes of brisk walking per week, accumulated in two different patterns, on cardiovascular risk. *Prev Med*, 41: 92-97.
- 35) Tully MA, Cupples ME, Chan WS, McGlade K, Young IS (2005): Brisk walking, fitness, and cardiovascular risk: a randomized controlled trial in primary care. *Prev Med*, 41: 622-628.
- 36) Audette JF, Jin YS, Newcomer R, Stein L, Duncan G, Frontera WR (2006): Tai Chi versus brisk walking in elderly women. *Age Ageing*, 35: 388-393.
- 37) Lee MR, Kim WS (2006): The effects of brisk walking versus brisk walking plus diet on triglycerides and apolipoprotein B levels in middle-aged overweight/obese women with high triglyceride levels. *Taehan Kanho Hakhoe Chi*, 36: 1352-1358.
- 38) Kirkwood L, Aldujaili E, Drummond S (2007): Effects of advice on dietary intake and/or physical activity on body composition, blood lipids and insulin resistance following a low-fat, sucrose-containing, high-carbohydrate, energy-restricted diet. *Int J Food Sci Nutr*, 58: 383-397.
- 39) Tully MA, Cupples ME, Hart ND, McEneny J, McGlade KJ, Chan WS, Young IS (2007): Randomised controlled trial of home-based walking programmes at and below current recommended levels of exercise in sedentary adults. *J Epidemiol Community Health*, 61: 778-783.
- 40) Coghill N, Cooper AR (2008): The effect of a home-based walking program on risk factors for coronary heart disease in hypercholesterolaemic men, a randomized controlled trial. *Prev Med*, 46: 545-551.
- 41) Pierce GL, Eskurza I, Walker AE, Fay TN, Seals DR (2011): Sex specific effects of habitual aerobic exercise on brachial artery flow-mediated dilation in middle-aged and older adults. *Clin Sci*, 120: 13-23.
- 42) Pagels P, Raustorp A, Archer T, Lidman U, Alricsson M (2012): Influence of moderate, daily physical activity on body composition and blood lipid profile in Swedish adults. *J Phys Act Health*, 9: 867-874.
- 43) Foulds HJ, Bredin SS, Charlesworth SA, Ivey AC, Warburton DER (2014): Exercise volume and intensity: a dose-response relationship with health benefit. *Eur J Appl Physiol*, 114: 1563-1571.
- 44) Kearney TM, Murphy MH, Davison GW, O'Kane MJ, Gallagher AM (2014): Accumulated brisk walking reduces arterial stiffness in overweight adults: evidence from a randomized control trial. *J Am Soc Hypertens*, 8: 117-126.
- 45) Johnson ST, Mundt C, Qiu W, Soprovich A, Wozniak L, Plotnikoff RC, Johnson JA (2015): Increase in daily steps after an exercise specialist led lifestyle intervention for adults with type 2 diabetes in primary care: a controlled implementation trial. *J Phys Act Health*, 12: 1492-1499.
- 46) Hui SS, Xie YJ, Woo J, Kwok TC (2016): Practicing Tai Chi had lower energy metabolism than walking but similar health benefits in terms of aerobic fitness, resting energy expenditure, body composition and self-perceived physical health. *Complement Ther Med*, 27: 43-50.

- 47) Zhang HJ, He J, Pan LL, Ma ZM, Han CK, Chen CS, Chen Z, Han HW, Chen S, Sun Q, Zhang JF, Li ZB, Yang SY, Li XJ, Li XY (2016): Effects of moderate and vigorous exercise on nonalcoholic fatty liver disease: a randomized clinical trial. *JAMA Intern Med*, 176: 1074-1082.
- 48) Blain H, Jaussent A, Picot MC, Maimoun L, Coste O, Masud T, Bousquet J, Bernard PL (2017): Effect of a 6-month brisk walking program on walking endurance in sedentary and physically deconditioned women aged 60 or older: a randomized trial. *J Nutr Health Aging*, 21: 1183-1189.
- 49) Blain H, Jaussent A, Picot MC, Çetinkaya C, Şişman AR, Çimrin D, Doğru Y, Kaya D (2017): The effect of 8-week different-intensity walking exercises on serum hepcidin, IL-6, and iron metabolism in pre-menopausal women. *Physiol Int*, 104: 52-63.
- 50) Herzig KH, Ahola R, Leppäluoto J, Jokelainen J, Jämsä T, Keinänen-Kiukaanniemi S (2014): Light physical activity determined by a motion sensor decreases insulin resistance, improves lipid homeostasis and reduces visceral fat in high-risk subjects: PreDiabEx study RCT. *Int J Obes*, 38: 1089-1096.
- 51) Ho SS, Dhaliwal SS, Hills AP, Pal S (2012): The effect of 12 weeks of aerobic, resistance or combination exercise training on cardiovascular risk factors in the overweight and obese in a randomized trial. *BMC Public Health*, 12: 1-10.
- 52) McNeilly AM, Davison GW, Murphy MH, Nadeem N, Trinick T, Duly E, Novials A, McEneny J (2011): Effect of α-lipoic acid and exercise training on cardiovascular disease risk in obesity with impaired glucose tolerance. *Lipids in Health and Disease*, 10: 1-9.
- 53) Mabire L, Mani R, Liu L, Mulligan H, Baxter D (2017): The influence of age, sex and body mass index on the effectiveness of brisk walking for obesity management in adults: a systematic review and meta-analysis. *J Phys Act Health*, 14: 389-407.
- 54) Nemoto K, Gen-no H, Masuki S, Okazaki K, Nose H (2007): Effects of high-intensity interval walking training on physical fitness and blood pressure in middle-aged and older people. *Mayo Clin Proc*, 82:803-11.
- 55) Donnelly JE, Blair SN, Jakicic JM, Manore MM, Rankin JW, Smith BK; American College of Sports Medicine (2009): American College of Sports and Medicine Position Stand: Appropriate physical activity intervention strategies for weight loss and prevention of weight regain for adults. *Med Sci Sports Exerc*, 41: 459 – 471.
- 56) Church TS, Earnest CP, Skinner JS, Blair SN (2007): Effects of different doses of physical activity on cardiorespiratory fitness among sedentary, overweight or obese postmenopausal women with elevated blood pressure. *JAMA*, 297: 2081 – 2091.
- 57) Donnelly JE, Hill JO, Jacobsen DJ, Pottenger J, Sullivan DK, Johnson SL, Heelan K, Hise M, Fennessey PV, Sonko B, Sharp T, Jakicic JM, Blair SN, Tran ZV, Mayo M, Gibson C, Washburn RA (2003): Effects of a 16-month randomized controlled exercise trial on body weight and composition in young, overweight men and women: the midwest exercise trial. *Arch Intern Med*, 163:1343 – 1350.
- 58) Durstine JL, Grandjean PW, Davis PG, Ferguson MA, Alderson NL, DuBose KD (2001): Blood lipid and lipoprotein adaptations to exercise: a quantitative analysis. *Sports Med*, 31:1033-1062.
- 59) Kelley GA, Kelley KS, Franklin B (2006): Aerobic exercise and lipids and lipoproteins in patients with cardiovascular disease: a meta-analysis of randomized controlled

- trials. *J Cardiopulm Rehabil*, 26:131-139.
- 60) Duncan JJ, Gordon NF, Scott CB (1991): Women walking for health and fitness: how much is enough? *JAMA*, 266:3295-3299.
- 61) Crouse SF, O'Brien BC, Grandjean PW, Lowe RC, Rohack JJ, Green JS, Tolson H (1997): Training intensity, blood lipids, and apolipoproteins in men with high cholesterol. *J Appl Physiol*, 82:270-277.